

## CHASELECTION

## Human IL28A/IFN-lambda 2, Tag Free

货号(Catalog Number): CY124FXXXX(L)

**别名(synonym):** interleukin-28A; IFN-lambda 2; IL28A; IL-28A; interferon, lambda 2

**来源(Source):** Human embryonic kidney cell, HEK293-derived human IL-28A/IFN-lambda 2 protein

**蛋白结构 (Structure):**

该蛋白不含标签

**基因 ID:** Q8IZJ0**氨基酸序列**


Val26-Val200

**分子量大小(MW)**


19.8 kDa (Monomer)

**纯度 (Purity)**

&gt; 95%, determined by SDS-PAGE

**SDS-PAGE**

4 ug/lane protein was resolved with SDS-PAGE under non-reducing (NR) and reducing (R) conditions and visualized by Coomassie Blue staining.

**Gel-filtration**

版本号: IN-PA-35-00

Size-exclusion chromatography of recombinant human IL28A protein (280 nm absorbance)

**内毒素含量 (Endotoxin)**


&lt;0.010 EU per 1 ug of the protein by the LAL method

**制剂(Formulation)**

Solution protein.

Dissolved in sterile PBS buffer.

This solution can be diluted into other aqueous buffers. Centrifuge the vial prior to opening.

**活性检测 (Biological Activity)**

Recombinant human IL28A antivirus activity of the HepG2 human hepatocellular carcinoma cells.

**储存与运输(Storage)**

Avoid repeated freeze-thaw cycles.

It is recommended that the protein be aliquoted for optimal storage.

36 months from date of receipt, -20 to -70 ° C as supplied.

**产品背景介绍 (Production)**

Interleukin-28A (IL-28A), IL-28B, and IL-29, also named interferon-lambda 2 (IFN-lambda 2), IFN-lambda 3, and IFN-lambda 1, respectively, are class II cytokine receptor ligands that are distantly related to members of the IL-10 family (11-13% aa sequence identity) and type I IFN family (15-19% aa sequence identity). The genes encoding these three cytokines are localized to chromosome 19 and each is composed of multiple exons. The exon organization of these genes is also found in the IL-10 family genes but

上海逐典生物科技有限公司

上海市自由贸易试验区李冰路 399 号抗体楼 5 楼 5018 室

www.chaselection.com

is distinct from the type I IFNs, which are encoded within a single exon. The expression of IL-28A,B, and IL-29 is induced by virus infection or double-stranded RNA. All three cytokines exert bioactivities that overlap those of type I IFNs, including antiviral activity and up-regulation of MHC class I antigen expression. The three proteins signal through the same heterodimeric receptor complex that is composed of the IL-10 receptor beta (IL-10 R beta ) and a novel IL-28 receptor alpha (IL-28 R alpha, also known as IFN-lambda R1). Ligand binding to the receptor complex induces Jak kinase activation and STAT1 and STAT2 tyrosine phosphorylation. The phosphorylated STAT1 and STAT2 complex with IFN-regulatory factor 9 (IRF-9) to form the IFN-stimulated regulatory factor 3 (ISGF-3) transcription factor complex that is translocated to the nucleus. ISGF-3 binds to the IFN-stimulated response element (ISRE) present in the regulatory regions of the target genes. Human IL-28A cDNA encodes a 200 amino acid (aa) residue precursor protein with a putative 25 aa signal peptide. It shares 94% and 67% aa sequence identity with human IL-28B and human IL-29,respectively.

