

CHASELECTION**Recombinant Human Noggin, Tag Free**

货号(Catalog Number): CY081FXXXX(L)

别名(synonym):NOG; Noggin; SYM1; symphalangism 1 (proximal);
synostoses (multiple) syndrome 1; SYNS1; SYNS1A**来源(Source):** Human embryonic kidney cell,
HEK293-derived human Noggin protein**蛋白结构 (Structure):**

该蛋白不含标签

基因 ID: Q13253**氨基酸序列:**


Gln28-Cys232

分子量大小(MW):

23.1 kDa[monomer]

纯度 (Purity):

> 95%, determined by SDS-PAGE

SDS-PAGE

2 ug/lane protein was resolved with SDS-PAGE under non-reducing (NR) and reducing (R) conditions and visualized by Coomassie Blue staining.


内毒素含量 (Endotoxin):

<0.010 EU per 1 ug of the protein by the LAL method.

制剂(Formulation):

Solution protein. Dissolved in sterile PBS buffer. This solution can be diluted into other aqueous buffers.

Centrifuge the vial prior to opening.

活性检测 (Biological Activity):

Recombinant human Noggin (Catalog # HF-2019) inhibits BMP-4 induced alkaline phosphatase production in the MC3T3E1 mouse preosteoblast cells.

储存与运输(Storage):

Avoid repeated freeze-thaw cycles. It is recommended that the protein be aliquoted for optimal storage.

36 months from date of receipt, -20 to -70 ° C as supplied.

Shipping with dry ice.

产品背景介绍 (Production):

Noggin is a secreted homodimeric glycoprotein that is an antagonist of bone morphogenetic proteins (BMPs). Human Noggin cDNA encodes a 232 amino acid (aa) precursor protein; cleavage of a 19 aa signal peptide generates the 213 aa mature protein which



contains an N-terminal acidic region, a central basic heparin-binding segment and a C-terminal cysteine-knot structure. Secreted Noggin probably remains close to the cell surface due to its binding of heparin-containing proteoglycans. Noggin is very highly conserved among vertebrates, such that mature human Noggin shares 99%, 99%, 98%, 97% and 89% aa sequence identity with mouse, rat, bovine, equine and chicken Noggin, respectively. Noggin binds some BMPs such as BMP-4 with high affinity and others such as BMP-7 with lower affinity. It antagonizes BMP bioactivities by blocking epitopes on BMPs that are needed for binding to both type I and type II receptors. During embryogenesis, Noggin antagonizes specific BMPs at defined times, for example, during neural tube, somite and cardiomyocyte growth and patterning. During skeletal development, Noggin prevents chondrocyte hyperplasia, thus allowing proper formation of joints. Mutations within the cysteine-knot region of human Noggin are linked to multiple types of skeletal dysplasias that result in apical joint fusions. Noggin is expressed in defined areas of the adult central nervous system and peripheral tissues such as lung, skeletal muscle and skin.

